

The Joint Commission

Journal on Quality and Patient Safety®

Improvement from
Front Office to Front Line

December 2010
Volume 36 Number 12



“Health systems that have done the hard work of developing a systematic approach to process improvement . . . will incorporate methodologic advances into their toolkit when it makes sense in the service of their overall process improvement activities.”

—VanKooy (p. 532)



Integrating Six Sigma and Lean Into the Improvement Toolkit

Features

Performance Improvement

- Optimizing the Process Improvement Toolkit
- Guiding Inpatient Quality Improvement: A Systematic Review of Lean and Six Sigma
- Pediatric HIVQUAL-T: Measuring and Improving the Quality of Pediatric HIV Care in Thailand, 2005–2007
- Kaiser Permanente’s Performance Improvement System, Part 2: Developing a Value Framework

Patient and Family Involvement

- Twelve Evidence-Based Principles for Implementing Self-Management Support in Primary Care

Department

Tool Tutorial

- Infection Preventionist Checklist to Improve Culture and Reduce Central Line–Associated Bloodstream Infections

Patient and Family Involvement

Twelve Evidence-Based Principles for Implementing Self-Management Support in Primary Care

Malcolm Battersby, Ph.D., M.B.B.S., F.R.A.N.Z.C.P., F.A.Ch.A.M.; Michael Von Korff, Ph.D.; Judith Schaefer, M.P.H.; Connie Davis, M.N., A.R.N.P.; Evette Ludman, Ph.D.; Sarah M. Greene, M.P.H.; Melissa Parkerton, M.A., A.B.S.; Edward H. Wagner, M.D., M.P.H., F.A.C.P.

In a rapidly aging population, two-thirds of persons over the age of 65 have multiple chronic conditions,¹ which has provoked urgent discussions of how society will provide for its health care needs.² The concept of an enabling society has been suggested as one way to enhance the abilities of citizens with chronic illness to successfully meet their own needs.³ Within the context of health care, *enablement* has been referred to as patient empowerment,⁴ self-management support (SMS),⁵ and collaborative care.⁶ The New Model of Care for family practice, for example, proposes a patient-centered team approach,⁷ which has been further elaborated into the concept of the patient-centered medical home, which includes SMS as one of its nine core components.⁸ Although the need to empower persons with chronic illness is apparent, it is less clear what specific steps health care providers can take to enhance self-management among their patients.

There is increasing evidence that effective self-management is essential to optimizing health outcomes for people with chronic conditions.^{5,6,9-13} This article seeks to identify approaches to SMS within routine clinical practice which can enable patients to successfully care for their chronic conditions.

The research team developed the Chronic Care Model (CCM)¹⁰ and, with the Institute of Healthcare Improvement (IHI), has been extensively involved in the IHI Breakthrough Series collaboratives' work with health care teams in attempting to implement the CCM—which includes SMS. Through a nominal group process, the authors used their extensive experience in clinical and quality improvement to identify key practices or processes that could guide clinicians or organizations in implementing SMS. We then used these elements to guide the literature review to identify interventions that could then be distilled into a set of principles.

We based our review on a definition that self-management consists of a set of tasks that includes developing knowledge of the condition(s) and treatment; medication management and

Article-at-a-Glance

Background: Recommendations to improve self-management support and health outcomes for people with chronic conditions in primary care settings are provided on the basis of expert opinion supported by evidence for practices and processes. Practices and processes that could improve self-management support in primary care were identified through a nominal group process. In a targeted search strategy, reviews and meta-analyses were then identified using terms from a wide range of chronic conditions and behavioral risk factors in combination with Self-Care, Self-Management, and Primary Care. On the basis of these reviews, evidence-based principles for self-management support were developed.

Findings: The evidence is organized within the framework of the Chronic Care Model. Evidence-based principles in 12 areas were associated with improved patient self-management and/or health outcomes: (1) brief targeted assessment, (2) evidence-based information to guide shared decision-making, (3) use of a nonjudgmental approach, (4) collaborative priority and goal setting, (5) collaborative problem solving, (6) self-management support by diverse providers, (7) self-management interventions delivered by diverse formats, (8) patient self-efficacy, (9) active follow-up, (10) guideline-based case management for selected patients, (11) linkages to evidence-based community programs, and (12) multifaceted interventions. A framework is provided for implementing these principles in three phases of the primary care visit: enhanced previsit assessment, a focused clinical encounter, and expanded postvisit options.

Conclusions: There is a growing evidence base for how self-management support for chronic conditions can be integrated into routine health care.

adherence; self-monitoring of disease and symptoms; managing the effects of illness on physical, emotional, and social role function; reducing health risks; preventative maintenance; and working collaboratively with health professionals.^{6,14} SMS includes actions by health care providers (formal and informal) that assist self-management.^{6,12} This definition assumes that optimal self-management is the product of a partnership between the patient, the family, and health care providers.

By including reduction of health risks and preventative maintenance of chronic conditions in the definition of self-management, we broadened the focus of the evidence from disease-specific self-management to alcohol, smoking, and related lifestyle domains, where much of the research on effective interventions to change behavior has been directed. We acknowledge the challenges in implementing SMS in routine practice and provide an example of how the principles may inform practice change by linking the principles to three phases of an enhanced primary care visit: (1) enhanced previsit assessment, (2) focused clinical encounter, and (3) expanded postvisit options. Finally, we comment on the need for more evidence of the cost-effectiveness of SMS interventions.

Methods

Literature Search. The set of practices or processes identified by the researchers was used to guide a literature search undertaken in October 2008–January 2009 to identify evidence from structured reviews and meta-analyses and summarized under key principles for implementing SMS in primary care. We focused the search strategy on evidence regarding a wide range of chronic conditions and behavioral risk factors. The English-language medical literature published between January 1980 and October 2008 was searched using the MEDLINE database of the National Library of Medicine and the Nursing and Allied Health database (from 1982). The Medical subject headings (MeSH) searched were Asthma, Diabetes, Coronary Arteriosclerosis, Heart Failure, Congestive, Depression, Pulmonary Disease, Chronic Obstructive, HIV/AIDS, Smoking Cessation, Alcohol-related Disorders, and Exercise, Self Care, and Primary Care. We also used a text word search of both databases using the term *self-management*. We limited the publication types to meta-analyses and systematic reviews. We also hand-searched all the systematic reviews of the Cochrane Effective Practice and Organization of Care (EPOC) group for those having relevance to SMS by affecting practice patterns.

Review of Relevance. The titles of the articles extracted by the search were reviewed for their relevance to the effectiveness of SMS. If potentially relevant, the full-text articles were

retrieved, and one author [J.S.] reviewed the studies for applicability.

Development of Thematic Content and Derivation of Principles. A second author [M.V.K.] reviewed the full-text articles, developed thematic content, and derived the 12 principles. The project team identified 123 reviews and meta-analyses, of which 83 were included. Evidence-based principles were chosen if they were supported by one or more reviews.

Findings

The 12 evidence-based principles are now provided.

Principle 1. Brief Targeted Assessment to Guide SMS: Assessment of clinical severity, functional status, patients' problems and goals, self-management behaviors, and barriers to self-management is integral to SMS. Direct evidence for assessment is sparse because it has not been isolated as an intervention. However, assessment of needs, preferences, behaviors, readiness, and barriers to self-management is a first step in all SMS interventions and assessment, and if done appropriately, is a way of engaging patients in their own care. It is not possible to match a patient with an intervention on the basis of preferences, readiness, or current behavior without assessment. These assessments then direct patient-specific behavior change interventions, collaborative care planning, and problem solving.¹⁰ A review of the effect of self-help materials for smoking cessation¹⁵ showed that interventions with assessment-based tailoring of materials to the individual were more effective than nontailored interventions. Strongest evidence of the value of assessment comes from the Public Health Service Tobacco Treatment Guideline,¹⁶ which found that assessment of smoking status (through a reminder) actually increased rates of clinician intervention and doubled cessation rates. Ebrahim et al.¹⁷ found that for secondary prevention programs following myocardial infarction, assessment and targeting interventions toward lifestyle risk factors resulted in significant benefits to patients.

Assessment of patient attributes such as readiness to change is integral to the use of motivational interviewing, an effective substance abuse intervention when used by clinicians.¹⁸ Katon et al. found that accurate diagnosis of comorbid depression and anxiety disorders in patients with chronic medical conditions was critical in the management of somatic symptoms.¹⁹ Similarly, Bower et al., in reviewing the treatment of depression in primary care, found that accurate diagnosis was a key predictor of treatment outcomes.²⁰

Principle 2. Information Alone Is Insufficient to Improve Patient Outcomes: Shared decision making should target evidence-based educational interventions that promote skill development. A review of adult asthma self-management education showed that information-only interventions did not have a significant effect²¹; “difficult asthma” psycho-education reduced hospitalizations in adults and children and improved symptoms in children.²² In a meta-analysis of diabetes self-management education, didactic patient education by itself was not sufficient to improve health outcomes.²³ In pediatric asthma, education, as compared with usual care, decreased the mean number of hospitalizations and emergency department visits, with more visits and interactive learning producing better outcomes.²⁴ Similarly, successful educational interventions for children with asthma incorporated skill-based training—either symptom-attack management or peak-flow training.²⁵ Warsi et al., who reviewed a range of self-management interventions, found evidence of reduced glycosylated hemoglobin (A1C) in diabetes, reduced systolic blood pressure in hypertension, and fewer attacks in asthma but nonsignificant changes in arthritis.²⁶ Interventions that involved face-to-face contact were associated with better outcomes. The authors noted that diabetes, hypertension, and asthma are conditions for which skills can be taught in diet, blood sugar control, and medication management, as opposed to arthritis, for which goals are less easy to define.

After a comprehensive clinical assessment, shared decision making between patient and clinician is used to determine which educational interventions are appropriate for the patient. Shared decision making is guided by evidence of what works. In smoking, the review conducted by Naughton et al. showed that for pregnant women, self-help interventions nearly doubled the quit rate, as compared with standard care.²⁷ However, self-help materials for smoking cessation have no additional benefit when used with advice from a health professional in adult smokers.^{15,22} In HIV/AIDS, patient support and education, delivered individually as opposed to in groups, improved adherence to antiretroviral therapy.²⁸

Principle 3. Use of a Nonjudgmental Approach: Clinicians more effectively support patient self-management when they provide evidence-based information with a nonjudgmental approach. Vogt et al. conducted a systematic review examining the attitudes of general practitioners (GPs) and family physicians toward smoking cessation.²⁹ A “sizeable minority” held negative beliefs about talking to their patients about smoking cessation, in the belief that it was “too time-consuming” (42%)

and ineffective (38%), and 22% lacked the confidence to discuss smoking with their patients. The authors concluded that interventions led by primary care physicians would be more effective if their own negative beliefs were addressed. Motivational interviewing is a directive counseling approach that explicitly avoids contradicting or judging a client.³⁰ Dunn et al. found that there was substantial evidence that motivational interviewing led to improved outcomes for patients with substance abuse and promising, yet inconclusive, evidence for its efficacy in HIV risk, diet/exercise, and smoking cessation.¹⁸ A meta-analysis of 72 studies revealed a significant effect for motivational interviewing for body mass index, total blood cholesterol, systolic blood pressure, and blood alcohol concentration but not for A1C or the number of cigarettes per day.³¹

Principle 4. Collaborative Priority and Goal Setting: Collaborative identification of priorities, goals, and specific plans for goal achievement improves self-management and outcomes. Locke et al. provide extensive evidence for what are considered the essential elements of goal setting and task motivation for improved outcomes.³² A review of interventions in chronic illness showed that collaborative problem definition, followed by setting realistic goals and developing a personalized care plan, led to improved outcomes.⁹ Harrington et al., who reviewed 20 studies that examined interventions aimed at improving communication and patient participation in medical consultations, found evidence that improvements in patients’ perceptions of the amount of control they have over their health, patients’ level of activity in maintaining their health, clinic attendance, and adherence to medical recommendations were associated with improved clinical outcomes.³³ In patients with congestive cardiac failure, comprehensive discharge planning and postdischarge support reduced hospital readmissions, reduced all-cause mortality, and improved quality of life.³⁴ A review of self-management education for adults with asthma identified improved outcomes, particularly when the interventions involved a written action plan, self-monitoring, and regular medical review.²¹ In chronic obstructive pulmonary disease (COPD), action plans helped people to recognize and act on an exacerbation of COPD symptoms.³⁵ For children and adolescents with asthma, there is limited evidence that a written plan alone produces better outcomes than no plan.³⁶ However, there is clear evidence that symptom-based plans are superior to peak flow-based plans.³⁷ In Type 2 diabetes, self-management education was found to improve A1C levels three months after the intervention.²³ The education programs included knowledge gain; lifestyle change; collabora-

tive goal-setting; and skill development, including coping skills.

Principle 5. Collaborative Problem Solving: Collaborative problem solving improves self-management and outcomes. Problem solving consists of a set of steps whereby individuals define a problem, brainstorm possible solutions, chose a strategy, try it, evaluate the response, and repeat the process. Problem solving therapy (PST) teaches patients how to use problem solving to identify strategies for achieving successful adoption of new behaviors and for overcoming barriers to change³⁸ and enables the translation of techniques for self-management into specific plans and behaviors.³⁹ A frequently cited intervention, the arthritis self-management program, includes problem-solving training.⁴⁰ This program resulted in improvements in pain of 20% compared with control groups, sustained for four years after the course. Lorig et al. emphasize that it is important to help people learn the skill of problem solving (as opposed to having a clinician do the problem solving for them) so they develop the capacity to address barriers themselves. A systematic review of problem solving in adults with diabetes showed that there was consistent evidence of an association between problem solving and A1C levels mediated through improvements in dietary behaviors.⁴¹ Problem solving was one of a combination of strategies in successful programs focusing on lifestyle change for diabetes.⁴² Better problem solving was associated with better self-care behavior but had little effect on diabetes control.⁴³

PST delivered by nurses in primary care improved depression outcomes for patients with depression.^{13,44} PST was effective in 72% of studies in a range of mental health problems in adults, children, and adolescents.⁴² Improved outcomes were associated with an increased number of sessions. In a meta-analytic review of coping, problem-focused coping correlated with improved health outcomes.⁴⁵ Support for problem solving was found in studies of self-management programs for arthritis.⁵

Principle 6. SMS by Diverse Providers: Diverse professionals and laypersons can effectively deliver SMS interventions if they have clearly defined tasks and roles and are trained to use evidence-based interventions. In a systematic review of diabetes education, Loveman et al. found that group programs achieved lifestyle change independent of which health professionals conducted the program as long as they were well trained.⁴⁶ A number of reviews have found improved health outcomes compared with controls when nurses or pharmacists were part of multifaceted interventions for diabetes,^{23,47} depres-

sion,¹³ and preventive services.⁴⁸ In a small number of trials of limited quality, dietitians were more effective than physicians in lowering blood cholesterol in the short to medium term, but there was no evidence that they were more effective than self-help resources or that they provided better outcomes than nurses.⁴⁹ Nurses have been key to successful interventions for depression in primary care,^{50,51} by providing telephone counseling for medication adherence, PST, and care management.⁴⁰ Thomas et al., who examined nurses' use of guidelines targeting management of hypertension, back pain, and hyperlipidemia, found that three of five studies reported improvements in processes of care and that six of another eight studies reported improvements in health outcomes.⁵² However, the authors advised caution in generalizing the findings because of methodologic problems in the studies. A systematic review of 18 studies of the use of community health workers in the care of patients with diabetes found positive effects of both lifestyle and self-care outcomes, as well as decreased "inappropriate health care use."⁵³ Foster et al. found that self-management interventions led by laypersons led to statistically significant improvement in patient confidence, health behaviors (exercise), small changes in pain, disability, fatigue, and depression but not quality of life.⁵⁴ A meta-analysis of studies that compared nurse practitioners to physicians in primary care found greater patient compliance with treatment recommendations, greater patient satisfaction, and resolution of pathological problems, for the nurse practitioners, although limitations of the studies reviewed were noted.⁵⁵

Principle 7. Self-Management Interventions Delivered by Diverse Formats: Self-management interventions can be effectively delivered via diverse modalities, including individual-, group-, telephone-, and self-instruction formats. Much of the evidence for the benefits of diverse modalities comes from the smoking-cessation literature. Group programs for smoking cessation were better than no intervention and equal to one session of individual counseling.⁵⁶ There was some extra but small benefit of group programs in addition to nicotine replacement but there was a wide variation in acceptance, with low uptake and more than a 30% dropout rate in some studies. Another review showed that telephone counseling using four to six phone calls led to higher quit rates than self-help materials alone, with no additional benefit from individual counseling or nicotine replacement.⁵⁷ Group training in self-management for people with Type 2 diabetes was effective in improving fasting blood glucose, A1C, knowledge, and need for diabetes medication.⁵⁸

Information technologies have emerged as innovative deliv-

ery systems with the potential to increase access to SMS. Kaltenhaler et al. found some evidence that computerized cognitive behavioral therapy (CBT) for anxiety, depression, and phobias was as effective as therapist-led CBT.⁵⁹ A review of randomized trials of information and communication technologies (ICTs) for chronic disease management did not show improvement in clinical outcomes except for cardiac diseases, which also showed lower mortality and health service utilization.⁶⁰ Systems for improving education and social support were effective. Bussey-Smith et al. found that computerized asthma education programs were effective in improving patient knowledge and symptoms but that clinical outcome effects were more varied.⁶¹ A systematic review of telemonitoring for patients with diabetes showed substantial decreases in A1C and complications and increases in patient self-efficacy, education, and receptiveness.⁶² More specifically, in a review of five studies of patients with diabetes, in which electronic monitoring was used to identify deficient medication compliance, adherence improved by 61%–79%.⁶³ Verhoven et al., who reviewed teleconsultation and videoconferencing in diabetes, found improved metabolic control and cost reduction but no change in quality of life.⁶⁴ A meta-analysis of 22 articles comparing Web-based and non-Web-based interventions for behavior change showed improvement in knowledge of nutritional status and asthma treatment, greater health care participation, and less health decline.⁶⁵

Principle 8. Patient Self-Efficacy: Enhancing patient self-efficacy regarding key chronic illness management tasks improves the process and outcomes of care. Self-efficacy is defined by Bandura as “people’s beliefs about their capabilities to produce designated levels of performance that exercise influence over events that affect their lives.”⁶⁶ Self-efficacy is not a process; it is an intermediate outcome or mediator of patient adoption of self-management behaviors and health behavior change. A review of fibromyalgia⁶⁷ notes the extensive work of Lorig and colleagues on self-management by patients with rheumatoid arthritis and osteoarthritis.^{68,69} Behavior and health status changes could be explained by patients’ attribution of improved control over symptoms. This sense of control is implicit in the definition of self-efficacy, that is, confidence in one’s ability to perform a task. The most powerful component of self-efficacy is mastery learning, which comes from achieving success in performing a new behavior. The arthritis self-management program, as cited earlier,⁴⁰ provides training in problem solving (Principle 5) and action planning, which are used to achieve a sense of mastery over new tasks. A meta-

analysis of psychological interventions for arthritis that featured combinations of relaxation, CBT, stress management, and coping skills training, either in groups or individually, found evidence that pain and disability improved compared with controls.⁷⁰ Significant effect sizes in coping and self-efficacy suggested that these factors may contribute to these outcomes. Asthma patients with enhanced self-efficacy (ability to self-adjust medications) showed more improvement in lung function than patients who relied on physicians to adjust their medication.³⁶ Similarly, in COPD, “self-sufficiency” in managing medical regimes was associated with a reduction in hospitalization.⁷¹ Self-efficacy includes self-prediction; a meta-analysis found that making a behavioral self-prediction significantly improved health-related behaviors.⁷²

Principle 9. Active Follow-Up: Ongoing follow-up, supported by feedback and reminders to both clinicians and patients, helps sustain self-management behaviors and improves patient outcomes. Most self-management interventions reported in the literature include some form of follow-up. Meta-analyses and reviews of multifaceted trials of diabetes,⁴⁹ depression,^{13,73,74} asthma,³⁶ and heart failure⁷⁵ have supported the role of reminders and follow-up to both patients and providers in improving health outcomes. Best outcomes in depression were achieved when follow-up was maintained for at least 12 months.⁷⁴ Two follow-up sessions to discuss medication substantially improved adherence, and nurse case management delivered over the telephone showed improved outcomes for depression.¹³ For heart failure, follow-up to specialized clinics or home visits was effective, whereas telephone follow-up to support primary care was not.⁷⁵ For diabetes, patient tracking systems for regular follow-up improved processes.⁴⁹ In reviewing studies assessing the effects of involving primary care physicians in routine review for complications of diabetes compared with specialist follow-up, Griffin et al. found that computerized recall for patients and their physicians could achieve standards of care as good as or better than outpatient care by specialists, at least in the short term.⁷⁶ Telemonitoring showed substantial decreases in A1C and increased patient self-efficacy,⁶⁴ and telecare, including nurses’ telephone follow-up of elderly people, improved clinical indicators for a range of chronic conditions.⁷⁷

Providing systematic follow-up is supported by evidence that follow-up increases rates of patient behavior change.^{78,79} Similar conclusions regarding the success of multiple strategies apply to the implementation of risk factor prevention in primary care settings^{50,80}; follow-up consisted of teleconference, newsletter, or individual instructions.⁵⁰ Overall, the reviewed

studies suggest that although reminders to attend appointments support self-management, follow-up's value depends on review of patient data, monitoring of progress toward goals, and use of problem solving to address barriers to achieving those goals.

Principle 10. Guideline-Based Case Management for Selected Patients: Case management can improve self-management and patient outcomes if (and only if) it is goal directed and guideline based. In reviewing case management, the dominant model of community mental health care, Marshall et al. noted its various definitions.⁸¹ These range from coordinating care to clinical case management (which is provided by a mental health professional using a therapeutic approach). In contrast, in assertive community treatment (ACT), a client is assigned to the team rather than an individual case worker.⁸¹ Marshall et al. found that in comparison to standard care, case management increased the number of community contacts and nearly doubled the number of admissions.⁸¹ Case management was no better on psychological, quality-of-life, or social measures. Marshall et al. recommended that community teams use ACT. Reviews of chronic care for depression describe case management as effective when it is targeted through behavioral approaches to selected cases, is goal-directed, enhances self-management,^{75,82} and provides direct feedback to the client and delivers psychological therapy.⁸³ Collaborative care for depression is effective when nurse case managers work with psychiatrists and primary care physicians.⁸⁴ A review of studies where the intervention was a specialist nurse in diabetes care alone, compared with usual care in hospital clinics or primary care, found that the nurse intervention may have improved A1C levels in the short term but that there was no difference from control groups during a 12-month period.⁸⁵ In addition, quality of life and emergency admissions were no different between groups. In minority ethnic groups with diabetes, case management improved A1Cs.⁸⁶ In another review, in which case management in primary care was defined as “a program that uses physicians or nonphysician providers to maintain continuous contact with patients via telephone or in-home visits in order to prevent disease exacerbation through intensive assessment and education techniques,”^{87(pp. 123–124)} only two of nine studies showed significant reduction in hospital admissions or bed days.⁸⁷ Both of the two studies examining clinical outcomes showed improvement, and six of the studies showed improvements in quality of life, patient satisfaction, and functional status. Interventions targeting specific diseases were more successful than general conditions supervised by generalists.

Principle 11. Linkages to Evidence-Based Community Programs: SMS should include community-based self-management programs that are evidence-based. Community-based telephone counseling for smoking cessation was shown to produce higher quit rates than usual care.⁸⁸ The Arthritis Self-Management Program (ASMP), developed by Lorig and colleagues at Stanford University^{89,90} (and cited earlier^{40,68,69}) is the most rigorously evaluated community-based program. This six-week, peer-led group program has produced gains in pain and fatigue management up to four years after the course.⁴⁰ A generic program for a range of chronic conditions based on the ASMP has produced improvements in pain and fatigue, as well as reductions in utilization and hospitalization at two-year follow-up.^{90,91} Newbold et al., in reviewing the ASMP and generic courses, documented a range of benefits from both programs and comparable outcomes when delivered by peers or health professionals.⁹² The use of ethnic link workers contributed to the benefits of diabetes management in ethnic minority groups,⁸⁸ and group interventions for chronic disease management held in culturally specific community settings had positive outcomes.⁵⁷

Principle 12. Multifaceted Interventions: Multifaceted interventions are more effective than single-component interventions. Effective SMS is typically provided by multifaceted rather than single-component interventions.^{6,11,12,44,93} Multifaceted interventions include patient registries that identify patients with specific chronic conditions⁹⁴ and linkage to appointment recall systems,⁹⁵ evidence-based guidelines, and outcome measures. In a review of trials of smoking cessation involving physicians and nonphysicians, Kottke et al. concluded that multifaceted interventions were more successful when teams of providers were involved and that combinations of group and individual sessions produced better outcomes than either alone.⁹⁶ Better outcomes were predicted by the number of interventions and the number of reinforcing sessions.⁹⁶

Discussion

We have provided a set of principles underpinned by evidence from randomized controlled trials to inform implementation of SMS in primary care. Evidence for the effectiveness of a number of these principles used in the Breakthrough Collaboratives has been described by Glasgow et al.¹¹ We acknowledge that these recommendations are most likely to be implemented when health care systems are able to afford to offer comprehensive programs for the patients who are willing and able to participate. The more comprehensive and intensive the programs,

Table 1. Integrating Self-Management Support (SMS) into Practice

Principles	Enhanced Visit Preparation	Focused Visit	Expand Options at Postvisit
1. Brief targeted assessment to guide SMS	●	●	—
2. Evidence-based information to guide shared decision making	—	●	●
3. Clinicians use a nonjudgmental approach	●	●	—
4. Collaborative priority and goal setting	●	●	—
5. Collaborative problem-solving	●	●	—
6. Diverse providers can offer SMS	—	—	●
7. Individual-, group-, telephone-, and self-instruction formats can be employed	—	—	●
8. Enhance patient self-efficacy	●	●	●
9. Ensure active follow-up	—	—	●
10. Guideline-based case management for selected patients	●	●	●
11. Linkages to evidence-based community-based self-management programs	—	—	●
12. Multifaceted interventions are more effective	●	●	●

the more likely they are to be effective in improving patient outcomes. In contrast, for disadvantaged populations, arguably the people who most need SMS, limited access may impede the implementation of the recommendations. In any case, embedding these principles into routine clinical practice is difficult; SMS is the least implemented of the six elements of the CCM.¹² Bodenheimer et al. documented the barriers to CCM, which included time, costs and the lack of reimbursement, inadequate information technology, and physician resistance to implementing chronic care management.⁹⁷ They recommended three redesign elements: (1) pre-activating patients before the clinic visit; (2) planned visits with a care manager who provides education and medical management to patients, either individually or in groups; and (3) sustained follow-up, either face to face, by telephone, or electronically provided⁹⁸ by a care team.⁹⁹ Glasgow et al.¹² provide a more detailed, sequential approach to integrating SMS, which is consistent with Bodenheimer, and expand the sustained follow-up to include linkages with community support and education programs.

Table 1 (above) shows how the 12 principles can be implemented via the three phases of patient care. By focusing on the enhanced assessment before the visit and on expanded post-visit options (for example, case management, self-management skills training, risk factor reduction), it may be possible to focus and strengthen SMS in the clinical encounter with only modest change in clinician behavior during the visit.

We have limited data on whether SMS interventions are cost-effective. Cost-effectiveness analyses are complex; the cost

per quality adjusted life years saved (QALYS) is the recommended outcome measure when comparing the cost-effectiveness of different interventions.¹⁰⁰ However, the impact on health outcomes with behavioral or self-management interventions may not be evident for many years, making QALYS and return on investment difficult to estimate. Although well-designed longitudinal studies should be conducted to measure the cost-effectiveness of self-management interventions, accurate cost analyses of interventions provides an intermediate step of benefit to both policy makers and funders. Readers are encouraged to collect their own cost data using feasible cost collection procedures.¹⁰¹

Conclusions

Health care is in the midst of a transition from expectations that disease will be managed primarily by physicians to a system that enables patients to effectively assume primary responsibility for managing chronic conditions. There is now a substantial evidence base that can be used to guide efforts to improve the abilities of health care teams to enable their patients to successfully manage chronic illness. The challenge before us is to integrate these evidence-based principles into routine patient care. **■**

This project was supported by Grant #46168, the Co-Management Learning Network, administered by the Health Research and Educational Trust of the American Hospital Association, and funded by The Robert Wood Johnson Foundation, grant #35678 Improving Chronic Illness Care from The Robert Wood Johnson Foundation to the MacColl Institute for Healthcare Innovation, Center for Health Studies, Group Health Cooperative, Seattle, and the Commonwealth Fund of New York Harkness Fellowship to Professor Malcolm Battersby.

Malcolm Battersby, Ph.D., M.B.B.S., F.R.A.N.Z.C.P., F.A.Ch.A.M., is Professor of Psychiatry and Director, Flinders Human Behaviour and Health Research Unit, Margaret Tobin Centre, Flinders University, Adelaide, Australia. **Michael Von Korff, Ph.D.**, is Senior Investigator, Center for Health Studies, Group Health Cooperative, Seattle, and **Judith Schaefer, M.P.H.**, is Research Associate. **Connie Davis, M.N., A.R.N.P.**, formerly Associate Director, Clinical Improvement, Improving Chronic Illness Care, Seattle, is a Gerontological Nurse Practitioner and Health Care Consultant, Hope, British Columbia, Canada. **Evette Ludman, Ph.D.**, is Senior Research Associate, Center for Health Studies; **Sarah M. Greene, M.P.H.**, is Research Associate; and **Melissa Parkerton, M.A., A.B.S.**, is Research Project Manager. **Edward H. Wagner, M.D., M.P.H., F.A.C.P.**, is Director, MacColl Institute, Center for Health Studies, Group Health Cooperative. Please address correspondence to Malcolm Battersby, malcolm.battersby@flinders.edu.au.

References

1. Wolff J.L., Starfield B., Anderson G.: Prevalence, expenditures, and complications of multiple chronic conditions in the elderly. *Arch Intern Med* 162:2269–2276, Nov. 11, 2002.
2. Hacker J.S.: *The Divided Welfare State: The Battle over Public and Private Social Benefits in the United States*. Cambridge, UK: Cambridge University Press 2002.
3. Gilbert N., Etzioni A.: *Transformation of the Welfare State: The Silent Surrender of Public Responsibility*. New York City: Oxford University Press, 2002.
4. Funnell M.M., Anderson R.M.: Patient empowerment: A look back, a look ahead. *Diabetes Educ* 29:454–464, May–Jun. 2003.
5. Bodenheimer T., et al.: Patient self-management of chronic disease in primary care. *JAMA* 288:2469–2475, Nov. 20, 2002.
6. Von Korff M., et al.: Collaborative management of chronic illness. *Ann Intern Med* 127:1097–1102, Dec. 15, 1997.
7. Martin J.C., et al.: The Future of Family Medicine: A collaborative project of the family medicine community. *Ann Fam Med* 2(suppl.1):S3–S32, Mar.–Apr. 2004.
8. American Academy of Family Physicians, et al.: *Joint Principles of the Patient Centered Medical Home*. Patient-Centered Primary Care Collaborative, Feb. 2007. <http://www.pccc.net/content/joint-principles-patient-centered-medical-home> (last accessed Nov. 4, 2010).
9. Wagner E.H., Austin B.T., Von Korff M.R.: Organizing care for patients with chronic illness. *Milbank Q* 74(41):511–544, 1996.
10. Wagner E.H., et al.: Improving chronic illness care: Translating evidence into action. *Health Aff (Millwood)* 20:64–78, Nov.–Dec. 2001.
11. Glasgow R.E., et al.: Self-management aspects of the Improving Chronic Illness Care Breakthrough Series: Implementation with diabetes and heart failure teams. *Ann Behav Med* 24:80–87, Spring 2002.
12. Glasgow R.E., et al.: Implementing practical interventions to support chronic illness self-management. *Jt Comm J Qual Saf* 29:563–574, Nov. 2003.
13. Gilbody S., et al.: Educational and organizational interventions to improve the management of depression in primary care: A systematic review. *JAMA* 289:3145–3151, Jun. 18, 2003.
14. Lorig K.R., Holman H.: Self-management education: History, definition, outcomes, and mechanisms. *Ann Behav Med* 26:1–7, Aug. 2003.
15. Lancaster T., Stead L.F.: Self-help interventions for smoking cessation. *Cochrane Database Syst Rev* (3):CD001118, Jul. 20, 2005.
16. U.S. Department of Health and Human Services: *Clinical Practice Guideline: Treating Tobacco Use and Dependence: 2008 Update*. Public Health Service, May 2008. http://www.surgeongeneral.gov/tobacco/treating_tobacco_use08.pdf (last accessed Nov. 4, 2010).
17. Ebrahim S., et al.: Multiple risk factor interventions for primary prevention of coronary heart disease. *Cochrane Database Syst Rev* (2):CD001561, 2000.
18. Dunn C., Deroo L., Rivara F.P.: The use of brief interventions adapted from motivational interviewing across behavioral domains: A systematic review. *Addiction* 96:1725–1742, Dec. 2001.
19. Katon W., Lin E.H., Kroenke K.: The association of depression and anxiety with medical symptom burden in patients with chronic medical illness. *Gen Hosp Psychiatry* 29:147–155, Mar.–Apr. 2007.
20. Bower P., et al.: Collaborative care for depression in primary care: Making sense of a complex intervention: Systematic review and meta-regression. *Br J Psychiatry* 189:484–493, Dec. 2006.
21. Gibson P.G., et al.: Limited (information only) patient education programs for adults with asthma. *Cochrane Database Syst Rev* (2):CD001005, 2002.
22. Smith J.R., et al.: A systematic review to examine the impact of psycho-educational interventions on health outcomes and costs in adults and children with difficult asthma. *Health Technol Assess* 9:iii–iv, 1–167, Jun. 2005.
23. Norris S.L., et al.: Self-management education for adults with Type 2 diabetes: A meta-analysis of the effect on glycemic control. *Diabetes Care* 25:1159–1171, Jul. 2002.
24. Coffman J.M., et al.: Effects of asthma education on children's use of acute care services: A meta-analysis. *Pediatrics* 121:575–586, Mar. 2008.
25. Wolf F.M., et al.: Educational interventions for asthma in children. *Cochrane Database Syst Rev* (1):CD000326, 2003.
26. Warsi A., et al.: Self-management education programs in chronic disease: A systematic review and methodological critique of the literature. *Arch Intern Med* 164:1641–1649, Aug. 9–23, 2004.
27. Naughton F., Prevost A.T., Sutton S.: Self-help smoking cessation interventions in pregnancy: A systematic review and meta-analysis. *Addiction* 103:566–579, Apr. 2008.
28. Rueda S., et al.: Patient support and education for promoting adherence to highly active antiretroviral therapy for HIV/AIDS. *Cochrane Database Syst Rev* (2):ACD001442, Jul. 19, 2006.
29. Vogt F., Hall S., Marteau T.M.: General practitioners' and family physicians' negative beliefs and attitudes towards discussing smoking cessation with patients: A systematic review. *Addiction* 100:1423–1431, Oct. 2005.
30. Miller W.R., Rollnick S.: *Motivational Interviewing: Preparing People to Change Addictive Behavior*. New York City: The Guilford Press, 1991.
31. Rubak S., et al.: Motivational interviewing: A systematic review and meta-analysis. *Br J Gen Pract* 55:305–312, Apr. 2005.
32. Locke E.A., Latham G.P.: Building a practically useful theory of goal setting and task motivation: A 35-year odyssey. *Am Psychol* 57:705–717, Sep. 2002.
33. Harrington J., Noble L.M., Newman S.P.: Improving patients' communication with doctors: A systematic review of intervention studies. *Patient Educ Couns* 52:7–16, Jan. 2004.
34. Phillips C.O., et al.: Comprehensive discharge planning with postdischarge support for older patients with congestive heart failure: A meta-analysis. *JAMA* 291:1358–1367, Mar. 17, 2004.
35. Turnock A.C., et al.: Action plans for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* (4):CD005074, Oct. 19, 2005.
36. Toelle B.G., Ram F.S.: Written individualised management plans for asthma in children and adults. *Cochrane Database Syst Rev* (2):CD002171, 2004.
37. Zemek R.L., Bhogal S.K., Ducharme F.M.: Systematic review of randomized controlled trials examining written action plans in children: What is the plan? *Arch Pediatr Adolesc Med* 162:157–163, Feb. 2008.
38. van den Hout J.H.C., Kerckhoffs-Hanssen M.R., Vlaeyen J.W.S.: The effect of problem solving therapy in mental health care and behavioral medi-

- cine: A systematic review. In van den Hout J.H.C.: *To Solve or Not Solve? Effects of Problems Solving Therapy and Graded Activity in Non-Specific Low Back Pain*. Maastricht The Netherlands: University of Maastricht, 2002, pp. 33–58.
<http://arno.unimaas.nl/show.cgi?did=10633> (last accessed Nov. 8, 2010).
39. Yates B.: *Self-Management: The Science and Art of Helping Yourself*. Belmont, CA: Wadsworth, 1985.
40. Lorig K.R., Mazonson P.D., Holman H.R.: Evidence suggesting that health education for self-management in patients with chronic arthritis has sustained health benefits while reducing health care costs. *Arthritis Rheum* 36:439–446, Apr. 1993.
41. Hill-Briggs F., Gemmill L.: Problem solving in diabetes self-management and control: A systematic review of the literature. *Diabetes Educ* 33:1032–1050, Nov.–Dec. 2007.
42. Whittmore R.: Strategies to facilitate lifestyle change associated with diabetes mellitus. *J Nurs Scholarsh* 32:225–232, 2000.
43. Hill-Briggs F.: Problem solving in diabetes self-management: A model of chronic illness self-management behavior. *Ann Behav Med* 25:182–193, Jun. 2003.
44. Bower P.: Primary care mental health workers: Models of working and evidence of effectiveness. *Br Journal Gen Pract* 52:926–933, Nov. 2002.
45. Penley J.A., Tomaka J., Wiebe J.S.: The association of coping to physical and psychological health outcomes: A meta-analytic review. *J Behav Med* 25:551–603, Dec. 2002.
46. Loveman E., et al.: The clinical and cost-effectiveness of patient education models for diabetes: A systematic review and economic evaluation. *Health Technol Assess* 7(22):iii, 1–190, 2003.
47. Renders C.M., et al.: Interventions to improve the management of diabetes mellitus in primary care, outpatient and community settings. *Cochrane Database Syst Rev* (1):CD001481, 2001.
48. Hulscher M.E., et al.: Interventions to implement prevention in primary care. *Cochrane Database Syst Rev* (1):CD000362, 2001.
49. Thompson R.L., et al.: Dietary advice given by a dietitian versus other health professional or self-help resources to reduce blood cholesterol. *Cochrane Database Syst Rev* (3):CD001366, 2003.
50. Simon G.E., et al.: Randomised trial of monitoring, feedback and management of care by telephone to improve treatment of depression in primary care. *BMJ* 320:550–554, Feb. 26, 2000.
51. Wagner E.H., Simon G.E.: Managing depression in primary care. *BMJ* 322:746–747, Mar. 31, 2001.
52. Thomas L., et al.: Guidelines in professions allied to medicine. *Cochrane Database Syst Rev* (2):CD000349, 2000.
53. Norris S.L., et al.: Effectiveness of community health workers in the care of persons with diabetes. *Diabet Med* 23:544–556, May 2006.
54. Foster G., et al.: Self-management education programmes by lay leaders for people with chronic conditions. *Cochrane Database Syst Rev* (4):CD005108, Oct. 17, 2007.
55. Brown S.A., Grimes D.E.: A meta-analysis of nurse practitioners and nurse midwives in primary care. *Nurs Res* 44:332–339, Nov.–Dec. 1995.
56. Stead L.F., Lancaster T.: Group behaviour therapy programmes for smoking cessation. *Cochrane Database Syst Rev* (2):CD001007, 2002.
57. Stead L.F., Perera R., Lancaster T.: Telephone counselling for smoking cessation. *Cochrane Database Syst Rev* (3):CD002850, Jul. 19, 2006.
58. Deakin T., et al.: Group based training for self-management strategies in people with type 2 diabetes mellitus. *Cochrane Database Syst Rev* (2):CD003417, Apr. 18, 2005.
59. Kaltenthaler E., et al.: Computerised cognitive behaviour therapy for depression and anxiety update: a systematic review and economic evaluation. *Health Technol Assess* 10:1–168, Sep. 2006.
60. Garcia-Lizana F., Sarria-Santamera A.: New technologies for chronic disease management and control: a systematic review. *J Telemed Telecare* 13(2):62–68, 2007.
61. Bussey-Smith K.L., Rossen R.D.: A systematic review of randomized control trials evaluating the effectiveness of interactive computerized asthma patient education programs. *Ann Allergy Asthma Immunol* 98:507–516, Jun. 2007.
62. Jaana M., Paré G.: Home telemonitoring of patients with diabetes: A systematic assessment of observed effects. *J Eval Clin Pract* 13:242–253, Apr. 2007.
63. Cramer J.A.: A systematic review of adherence with medications for diabetes. *Diabetes Care* 27:1218–1224, May 2004.
64. Verhoeven F., et al.: The contribution of teleconsultation and videoconferencing to diabetes care: A systematic literature review. *J Med Internet Res* 9:e37, Dec. 14, 2007.
65. Wantland D.J., et al.: The effectiveness of Web-based vs. non-Web-based interventions: A meta-analysis of behavioral change outcomes. *J Med Internet Res* 6:, Nov. 10, 2004. <http://www.jmir.org/2004/4/e40/> (last accessed Nov. 4, 2010).
66. Bandura A.: Self-efficacy. In V.S. Ramachandran (ed.): *Encyclopedia of Human Behavior*, vol. 4, pp. 71–81. New York City: Academic Press, pp. 71–81. (Reprinted in H. Friedman ed.: *Encyclopedia of Mental Health*. San Diego: Academic Press, 1998). <http://www.des.emory.edu/mfp/BanEncy.html#top> (last accessed Nov. 4, 2010).
67. Burckhardt C.S., Bjelle A.: Education programmes for fibromyalgia patients: description and evaluation. *Baillieres Clin Rheumatol*. 8(4):935–55, November 1994.
68. Lorig K., Holman H.: Arthritis Self-Management Studies: A Twelve-Year Review. *Health Education Quarterly* 20(1):17–28, Spring 1993.
69. Lorig K., et al.: The beneficial outcomes of the arthritis self-management course are not adequately explained by behavior change. *Arthritis Rheum* 32:91–95, Jan. 1989.
70. Astin J.A., et al.: Psychological interventions for rheumatoid arthritis: a meta-analysis of randomized controlled trials. *Arthritis Rheum* 47:291–302, Jun. 2002.
71. Effing T.W., et al.: Self-management education for patients with chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2007 (3):CD002990, Oct. 17, 2007.
72. Sprott D.E., et al.: Self-prediction and patient health: influencing health-related behaviors through self-prophecy. *Med Sci Monit* 12:RA85–RA91, May 2006.
73. Von Korff M., Goldberg D.: Improving outcomes in depression. *BMJ* 323(7439):948–949, Oct. 27, 2001.
74. Ellis P.M., Smith D.A., Beyond Blue: The national depression initiative. Treating depression: The Beyond Blue Guidelines for Treating Depression in Primary Care. “Not so much what you do but that you keep doing it.” *Med J Aust* 176 Suppl:S77–S83(suppl.), May 20, 2002.
75. McAlister F.A., et al.: A systematic review of randomized trials of disease management programs in heart failure. *Am J Med* 110:378–384, Apr. 2001.
76. Griffin S., Kinmonth A.L.: Systems for routine surveillance for people with diabetes mellitus. *Cochrane Database Syst Rev* (1):CD000541, Jan. 21, 2009.
77. Barlow J., et al.: A systematic review of the benefits of home telecare for frail elderly people and those with long-term conditions. *J Telemed Telecare* 13:172–179, 2007.
78. Lancaster T., Silagy C., Fowler G.: Training health professionals in smoking cessation. *Cochrane Database Syst Rev* 2000(3):CD000214, 2000.
79. Silagy C., Stead L.F.: Physician advice for smoking cessation. *Cochrane Database Syst Rev* (4):CD000165, 2001.
80. Glasgow R.E., et al.: Translating what we have learned into practice: Principles and hypotheses for interventions addressing multiple behaviors in primary care. *Am J Prev Med* 27(Suppl.):88–101, Aug. 2004
81. Marshall M., et al.: Case management for people with severe mental disorders. *Cochrane Database Syst Rev* (2):CD000050, 2000.
82. Gensichen J., et al.: Case management to improve major depression in pri-

- mary health care: a systematic review. *Psychol Med* 36:7–14, Aug. 2006.
83. Christensen H., et al.: Models in the delivery of depression care: A systematic review of randomised and controlled intervention trials. *BMC Fam Pract* 9:25, May 5, 2008.
84. Gilbody S., et al.: Collaborative care for depression: A cumulative meta-analysis and review of longer-term outcomes. *Arch Intern Med* 166:2314–2321, Nov. 27, 2006.
85. Loveman E., Royle P., Waugh N.: Specialist nurses in diabetes mellitus. *Cochrane Database Syst Rev* (2):CD003286, 2003.
86. Saxena S., et al.: Systematic review of primary healthcare interventions to improve diabetes outcomes in minority ethnic groups. *J Ambul Care Manage* 30:218–230, Jul.–Sep. 2007.
87. Ferguson J.A., Weinberger M.: Case management programs in primary care. *J Gen Intern Med* 13:123–126, Feb. 1998.
88. Stead L.F., Lancaster T., Perera R.: Telephone counselling for smoking cessation. *Cochrane Database Syst Rev* (1):Jul. 19, 2006.
89. Lorig K.R., et al.: Outcomes of self-help education for patients with arthritis. *Arthritis Rheum* 28:680–685, Jun. 1985.
90. Lorig K.R., et al.: Evidence suggesting that a chronic disease self-management program can improve health status while reducing hospitalization: A randomized trial. *Med Care* 37:5–14, Jan. 1999.
91. Lorig K.R., et al.: Effect of a self-management program on patients with chronic disease. *Eff Clin Pract* 4:256–262, Nov.–Dec. 2001.
92. Newbould J., Taylor D., Bury M.: Lay-led self-management in chronic illness: A review of the evidence. *Chronic Illn* 2:249–261, Dec. 2006.
93. Stone A.A., Porter L.S.: Psychological coping: Its importance for treating medical problems. *Mind/Body Med* 1:46–54, 1995.
94. Haynes R.B., et al.: Interventions for helping patients to follow prescriptions for medications. *Cochrane Database Syst Rev* (2):CD000011, 2002.
95. Stone E.G., et al.: Interventions that increase use of adult immunization and cancer screening services: A meta-analysis. *Ann Intern Med* 136:641–651, May 7, 2002.
96. Kottke T.E., et al.: Attributes of successful smoking cessation interventions in medical practice: A meta-analysis of 39 controlled trials. *JAMA* 259:2883–2889, May 20, 1988.
97. Bodenheimer T., et al.: What are the facilitators and barriers in physician organizations' use of care management processes? *Jt Comm J Qual Saf* 30:505–514, Sep. 2004.
98. Bodenheimer T.: Helping patients improve their health-related behaviors: what system changes do we need? *Dis Manag* 8:319–330, Oct. 2005.
99. Bodenheimer T., Laing B.Y.: The teamlet model of primary care. *Ann Fam Med* 5:457–461, 2007.
100. Torrance G.W.: Measurement of health state utilities for economic appraisal. *J Health Econ* 5:1–30, Mar. 1986.
101. Ritzwoller D., et al.: Costing behavioral interventions: A practical guide to enhance translation. *Ann Behav Med* 37:218–227, Apr. 2009.



VA Patient Safety Fellowship

The VA National Center for Patient Safety, in cooperation with the VA Office of Academic Affiliations is pleased to offer a fellowship opportunity in Patient Safety beginning July 2011. The fellows are based at one of seven VA Medical Center sites: Ann Arbor, MI; Indianapolis, IN; Lexington, KY; Philadelphia, PA; Pittsburgh, PA; Tampa, FL; and White River Junction, VT. Open to any healthcare professional with a Master's degree or higher (such as physicians, nurses, psychologists, pharmacists, social workers and health care administrators). The one-year fellowship is designed to develop the next generation of leaders in patient safety. Fellows receive intensive training in clinical aspects of patient safety, health services research methodology, and leadership.

Start or rejuvenate your career. Join the VA for a year of mentoring and professional development. Additional information and fellowship requirements can be found at www.patientsafety.gov/jobs.html.

To apply, send a cover letter and CV to the Program Director(s) at the sites of interest.

Ann Arbor, MI	Linda Williams	Linda.Williams7@va.gov
Indianapolis, IN	Richard Frankel	Richard.Frankel@va.gov
Lexington, KY	Joe Conigliaro	Joseph.Conigliaro@va.gov
Philadelphia, PA	Gregg Lipschik	Gregg.Lipschik@va.gov
Pittsburgh, PA	David Eibling	David.Eibling@va.gov
Tampa, FL	Tatjana Bulat	Tatjana.Bulat@va.gov
White River Jct., VT	Vince Watts	Bradley.Watts@va.gov

EOE



PHYSICIAN EDUCATOR MEDICAL TEAM TRAINING (MTT)

The VA National Center for Patient Safety (NCPS), located in Ann Arbor, Michigan, is soliciting applications for the position of Physician Educator with the Medical Team Training (MTT)

Program. VHA is the largest integrated healthcare delivery system in the nation. NCPS is an internationally recognized center that develops and implements patient safety policies, tools, education, and information systems throughout the VA healthcare system (www.patientsafety.gov).

The MTT Program is designed to improve outcomes of patient care and staff job satisfaction by implementing Crew Resource Management communication tools from aviation with briefings and debriefings in the clinical workplace. MTT is provided to VA Medical Centers (VAMC) nationwide to improve patient care by enhancing communication in the delivery of care by healthcare professionals. The Physician Educator collaborates with multiple disciplines in the assessment, planning, organization, implementation, and evaluation of an effective MTT educational program, and supports other NCPS clinical education and training programs. This includes development of educational tools critical to the program such as teaching films, cognitive aids, and clinical scenarios for role play. The Physician Educator executes position responsibilities that demonstrate leadership, experience, and creative approaches to improvement of complex care delivery.

Board certified/board eligible physicians with a current, unrestricted license to practice medicine in any of the 50 U.S. states may apply. Previous experience and strong skills and ability in public speaking/teaching are required. The preferred candidate will have completed formal training in the principles of adult education and instructional techniques; formal training in patient safety; and will have experience teaching in a simulation environment. Frequent overnight travel (9-12 nights/month) is required. Must be U.S. citizen.

For additional information and application instructions, see www.patientsafety.gov/jobs.html

